

CS145: INTRODUCTION TO DATA MINING

3: Vector Data: Logistic Regression

Instructor: Yizhou Sun


yzsun@cs.ucla.edu

October 9, 2017

Methods to Learn

	Vector Data	Set Data	Sequence Data	Text Data
Classification	Logistic Regression ; Decision Tree; KNN SVM; NN			Naïve Bayes for Text
Clustering	K-means; hierarchical clustering; DBSCAN; Mixture Models			PLSA
Prediction	Linear Regression GLM*			
Frequent Pattern Mining		Apriori; FP growth	GSP; PrefixSpan	
Similarity Search			DTW	

Vector Data: Logistic Regression

- Classification: Basic Concepts 
- Logistic Regression Model
- Generalized Linear Model*
- Summary

Supervised vs. Unsupervised Learning

- Supervised learning (classification)
 - Supervision: The training data (observations, measurements, etc.) are accompanied by **labels** indicating the class of the observations
 - New data is classified based on the training set
- Unsupervised learning (clustering)
 - The class labels of training data is unknown
 - Given a set of measurements, observations, etc. with the aim of establishing the existence of classes or clusters in the data

Prediction Problems: Classification vs. Numeric Prediction

- Classification
 - predicts categorical class labels
 - classifies data (constructs a model) based on the training set and the values (**class labels**) in a classifying attribute and uses it in classifying new data
- Numeric Prediction
 - models continuous-valued functions, i.e., predicts unknown or missing values
- Typical applications
 - Credit/loan approval:
 - Medical diagnosis: if a tumor is cancerous or benign
 - Fraud detection: if a transaction is fraudulent
 - Web page categorization: which category it is


Classification—A Two-Step Process (1)

- **Model construction**: describing a set of predetermined classes
 - Each tuple/sample is assumed to belong to a predefined class, as determined by the **class label attribute**
 - For data point i : $\langle x_i, y_i \rangle$
 - Features: x_i ; class label: y_i
 - The model is represented as classification rules, decision trees, or mathematical formulae
 - Also called classifier
- The set of tuples used for model construction is **training set**


Classification—A Two-Step Process (2)

- **Model usage**: for classifying future or unknown objects
- **Estimate accuracy of the model**
 - The known label of test sample is compared with the classified result from the model
 - **Test set** is independent of training set (otherwise overfitting)
 - **Accuracy** rate is the percentage of test set samples that are correctly classified by the model
 - Most used for binary classes
- **If the accuracy is acceptable, use the model to classify new data**
- Note: If *the test set* is used to select models, it is called **validation (test) set**


Process (1): Model Construction



Process (2): Using the Model in Prediction



Vector Data: Logistic Regression

- Classification: Basic Concepts
- Logistic Regression Model 
- Generalized Linear Model*
- Summary


Linear Regression VS. Logistic Regression

- Linear Regression (prediction)
 - Y : *continuous value* $(-\infty, +\infty)$
 - $Y = \mathbf{x}^T \boldsymbol{\beta} = \beta_0 + x_1\beta_1 + x_2\beta_2 + \cdots + x_p\beta_p$
 - $Y|\mathbf{x}, \boldsymbol{\beta} \sim N(\mathbf{x}^T \boldsymbol{\beta}, \sigma^2)$
- Logistic Regression (classification)
 - Y : *discrete value from m classes*
 - $p(Y = C_j) \in [0,1]$ and $\sum_j p(Y = C_j) = 1$

Logistic Function

- Logistic Function / sigmoid function:

$$\sigma(x) = \frac{1}{1+e^{-x}}$$



Note: $\sigma'(x) = \sigma(x)(1 - \sigma(x))$

Modeling Probabilities of Two Classes


- $P(Y = 1|X, \beta) = \sigma(X^T \beta) = \frac{1}{1 + \exp\{-X^T \beta\}} = \frac{\exp\{X^T \beta\}}{1 + \exp\{X^T \beta\}}$
- $P(Y = 0|X, \beta) = 1 - \sigma(X^T \beta) = \frac{\exp\{-X^T \beta\}}{1 + \exp\{-X^T \beta\}} = \frac{1}{1 + \exp\{X^T \beta\}}$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix}$$

- In other words
 - $Y|X, \beta \sim \text{Bernoulli}(\sigma(X^T \beta))$


The 1-d Situation

- $P(Y = 1|x, \beta_0, \beta_1) = \sigma(\beta_1 x + \beta_0)$



Example

Regression of Sex on Height



Q: What is β_0 here?

Parameter Estimation

- MLE estimation
 - Given a dataset D , with n data points
 - For a single data object with attributes \mathbf{x}_i , class label y_i
 - Let $p_i = p(Y = 1|\mathbf{x}_i, \beta)$, the prob. of i in class 1
 - The probability of observing y_i would be
 - If $y_i = 1$, then p_i
 - If $y_i = 0$, then $1 - p_i$
 - Combining the two cases: $p_i^{y_i}(1 - p_i)^{1-y_i}$

$$L = \prod_i p_i^{y_i} (1 - p_i)^{1-y_i} = \prod_i \left(\frac{\exp\{X^T \beta\}}{1 + \exp\{X^T \beta\}} \right)^{y_i} \left(\frac{1}{1 + \exp\{X^T \beta\}} \right)^{1-y_i}$$

Optimization

- Equivalent to maximize log likelihood
- $L = \sum_i y_i \mathbf{x}_i^T \beta - \log(1 + \exp\{\mathbf{x}_i^T \beta\})$
- Gradient ascent update:

- $$\beta^{new} = \beta^{old} + \boxed{\eta} \frac{\partial L(\beta)}{\partial \beta}$$

Step size

- Newton-Raphson update

- $$\beta^{new} = \beta^{old} - \left(\frac{\partial^2 L(\beta)}{\partial \beta \partial \beta^T} \right)^{-1} \frac{\partial L(\beta)}{\partial \beta}$$

- where derivatives are evaluated at β^{old}

First Derivative

$$\begin{aligned}\frac{\partial L(\beta)}{\beta_{1j}} &= \sum_{i=1}^N y_i x_{ij} - \sum_{i=1}^N \frac{x_{ij} e^{\beta^T x_i}}{1 + e^{\beta^T x_i}} \\ &= \sum_{i=1}^N y_i x_{ij} - \sum_{i=1}^N p(x_i; \beta) x_{ij} \\ &= \sum_{i=1}^N x_{ij} (y_i - p(x_i; \beta))\end{aligned}$$

$p(x_i; \beta)$

$$j = 0, 1, \dots, p$$

Second Derivative

- It is a $(p+1)$ by $(p+1)$ matrix, Hessian Matrix, with j th row and n th column as

$$\begin{aligned} & \frac{\partial L(\beta)}{\partial \beta_{1j} \partial \beta_{1n}} \\ = & - \sum_{i=1}^N \frac{(1 + e^{\beta^T x_i}) e^{\beta^T x_i} x_{ij} x_{in} - (e^{\beta^T x_i})^2 x_{ij} x_{in}}{(1 + e^{\beta^T x_i})^2} \\ = & - \sum_{i=1}^N x_{ij} x_{in} p(x_i; \beta) - x_{ij} x_{in} p(x_i; \beta)^2 \\ = & - \sum_{i=1}^N x_{ij} x_{in} p(x_i; \beta) (1 - p(x_i; \beta)) . \end{aligned}$$


What about Multiclass Classification?

- It is easy to handle under logistic regression, say M classes

- $$P(Y = j|X) = \frac{\exp\{X^T \beta_j\}}{1 + \sum_{m=1}^{M-1} \exp\{X^T \beta_m\}}, \text{ for } j = 1, \dots, M-1$$

- $$P(Y = M|X) = \frac{1}{1 + \sum_{m=1}^{M-1} \exp\{X^T \beta_m\}}$$

Vector Data: Logistic Regression

- Classification: Basic Concepts
- Logistic Regression Model
- Generalized Linear Model* 
- Summary

Recall Linear Regression and Logistic Regression

- Linear Regression
 - $y|\mathbf{x}, \beta \sim N(\mathbf{x}^T \beta, \sigma^2)$
- Logistic Regression
 - $y|\mathbf{x}, \beta \sim \text{Bernoulli}(\sigma(\mathbf{x}^T \beta))$
- How about other distributions?
 - Yes, as long as they belong to exponential family

Exponential Family

- Canonical Form

- $p(y; \eta) = b(y) \exp(\eta^T T(y) - a(\eta))$

- η : natural parameter

- $T(y)$: sufficient statistic

- $a(\eta)$: log partition function for normalization

- $b(y)$: function that only dependent on y

Examples of Exponential Family

- Many:

$$p(y; \eta) = b(y) \exp(\eta^T T(y) - a(\eta))$$

- Gaussian, Bernoulli, Poisson, beta, Dirichlet, categorical, ...

- For Gaussian (not interested in σ)

$$\begin{aligned} p(y; \mu) &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}(y - \mu)^2\right) \\ &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}y^2\right) \cdot \exp\left(\mu y - \frac{1}{2}\mu^2\right) \end{aligned}$$

$$\begin{aligned} \eta &= \mu \\ T(y) &= y \\ a(\eta) &= \mu^2/2 \\ &= \eta^2/2 \\ b(y) &= (1/\sqrt{2\pi}) \exp(-y^2/2) \end{aligned}$$

- For Bernoulli


$$\begin{aligned} p(y; \phi) &= \phi^y (1 - \phi)^{1-y} \\ &= \exp(y \log \phi + (1 - y) \log(1 - \phi)) \\ &= \exp\left(\underbrace{\left(\log\left(\frac{\phi}{1 - \phi}\right)\right)}_{\eta} y + \log(1 - \phi)\right) \end{aligned}$$

$$\begin{aligned} T(y) &= y \\ a(\eta) &= -\log(1 - \phi) \\ &= \log(1 + e^\eta) \\ b(y) &= 1 \end{aligned}$$

Recipe of GLMs

- Determines a distribution for y
 - E.g., Gaussian, Bernoulli, Poisson
- Form the linear predictor for η
 - $\eta = \mathbf{x}^T \boldsymbol{\beta}$
- Determines a link function: $\mu = g^{-1}(\eta)$
 - Connects the linear predictor to the mean of the distribution
 - E.g., $\mu = \eta$ for Gaussian, $\mu = \sigma(\eta)$ for Bernoulli, $\mu = \exp(\eta)$ for Poisson

Vector Data: Logistic Regression

- Classification: Basic Concepts
- Logistic Regression Model
- Generalized Linear Model*
- Summary 

Summary

- What is classification
 - Supervised learning vs. unsupervised learning, classification vs. prediction
- Logistic regression
 - Sigmoid function, multiclass classification
- Generalized linear model*
 - Exponential family, link function